

Germes and Sylows for structure group of solutions to the Yang–Baxter equation

GRYB23 Conference

Edouard Feingessicht (LMNO, Caen, France)

June 2023, Blankenberge

Yang–Baxter Equation

Set-theoretical solution of the YBE (*Drinfeld '92*)

(X, r) where X is a set and $r : X \times X \rightarrow X \times X$ a bijection, such that

$$r_1 r_2 r_1 = r_2 r_1 r_2$$

where $r_i : X \times X \times X \rightarrow X \times X \times X$ acts on the coordinates i and $i + 1$.

For any X , $r(x, y) = (y, x)$ defines a solution.

Definition (*Etingof–Schedler–Soloviev '99*)

Denote $r(x, y) = (\lambda(x, y), \rho(x, y))$. (X, r) is said to be:

- Involutive if $r^2 = \text{id}_{X \times X}$
- Left non-degenerate (resp. right) if $\lambda(x, -)$ (resp. $\rho(-, y)$) is a bijection for any x (resp. y).

Cycle sets

Cycle set (Rump '05)

$(S, *)$ where S is a set and $*$ a binary operation such that for any s in S the map $\psi(s) : t \mapsto s * t$ is bijective, and for all s, t, u in S

$$(s * t) * (s * u) = (t * s) * (t * u).$$

Example: $S = \{s_1, \dots, s_n\}$ and $s * s_i = s_{\sigma(i)}$, with $\sigma = (12 \dots n)$
($\psi(s_i) = \sigma$).

Theorem (Rump '05)

There is a bijective correspondence in the finite cases

involutive left non-degenerate solutions \longleftrightarrow Cycle sets

Structure groups

Definition-Proposition (Etingof–Schedler–Soloviev '99, Rump '05)

Define the structure group G (resp. monoid M) by the presentation

$$\langle X \mid xy = x'y' \text{ if } r(x, y) = (x', y') \rangle \longleftrightarrow \langle S \mid s(st) = t(ts) \rangle.$$

Example: $S = \{s_1, s_2\}$ with $\psi(s_i) = (12)$ yields $M = \langle s_1, s_2 \mid s_1^2 = s_2^2 \rangle^+$.

Suppose S finite and fix an enumeration $S = \{s_1, \dots, s_n\}$.

Representation (Dehornoy '15)

We define the morphism $\Theta : G \rightarrow \mathrm{GL}_n(\mathbb{Q}[q, q^{-1}])$ induced by

$$s_i \mapsto D_i P_{s_i} = \mathrm{diag}(1, \dots, q, \dots, 1) \cdot P_{\psi(s_i)}.$$

Example: $\Theta(s_1) = \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & q \\ 1 & 0 \end{pmatrix}$ et $\Theta(s_2) = \begin{pmatrix} 0 & 1 \\ q & 0 \end{pmatrix}$.

- A monomial matrix m decomposes uniquely as $m = D_m P_m = P_m D'_m$.

I-Structure

Theorem (I-structure) (Gateva-Ivanova–Van den Bergh '98)

The only permutation matrix in $\Theta(G)$ is the identity.

- In other words, $\Theta(f)$ is uniquely determined by $D_{\Theta(f)}$ for f in G .

Theorem (Dehornoy '15)

Θ is a faithful representation.

- $G < \mathbb{Z}^n \rtimes \mathfrak{S}_n$ such that projecting on the 1st coordinate is bijective.

Theorem (Chouraqui '10)

G is a Garside group.

- G has a nice "lattice" structure with a preferred element Δ
- B_n are Garside groups, they can be "recovered" from \mathfrak{S}_n

Dehornoy's class and germ

$s_i^{[k]}$ the unique element of G with diagonal part D_i^k .

$$s_2^{[3]} = \begin{pmatrix} 0 & 1 \\ q^3 & 0 \end{pmatrix}$$

Proposition (Dehornoy's class)

There exists $d \in \mathbb{N}$ such that $s^{[d]}$ is diagonal for all $s \in S$.

Example: If $S = \{s_1, \dots, s_n\}$ et $\psi(s_i) = (12 \dots n)$, $d = n$.

Theorem (Germ) (Dehornoy '15)

(M, Δ^{d-1}) can be "recovered" from $\overline{G} = G / \langle s^{[d]} \rangle$ finite.

- $G \twoheadrightarrow \overline{G}$ amounts to evaluating $q = \exp(\frac{2i\pi}{d})$.

A conjecture on the class

Using Vendramin's enumeration :

n	$d_{\max}(n)$
3	3
4	4
5	6
6	8
7	12
8	15
9	24
10	30

Maximum of the classes of cycle sets with size n

Conjecture (F.)

$d_{\max}(n)$ is equal to "Maximum of products of distinct partitions of n ".

- Example: $n = 9 = 2 + 3 + 4$, and $2 \cdot 3 \cdot 4 = 24$ is maximal.
- A034893 on the OEIS. And Došlić gave an explicit formula (with T_m).

More on Dehornoy's class

Denote $\mathcal{G} < \mathfrak{S}_n$ the group generated by the $\psi(s)$'s.

Proposition

If $T : s \mapsto s * s$, we have : $o(T) \mid d \mid \#\mathcal{G} \mid d^n$.

In particular, d and $\#\mathcal{G}$ have the same prime divisors.

Proposition (F.)

If $s * s = s$ for all s and \mathcal{G} is abelian, the conjecture is verified.

Remark : It is "enough" to classify braces with additive group $(\mathbb{Z}/d\mathbb{Z})^n$, $d \leq d_{\max}(n)$.

Sylow for the germs

Theorem (*Lebed-Ramírez-Vendramin '22, F.*)

$G^{[k]} = \langle s^{[k]} \rangle$ induces a cycle set structure on $S^{[k]} = \{s^{[k]}\}_{s \in S}$.

Moreover, its class is $\frac{d}{d \wedge k}$ (if $k \leq d$).

- $G^{[k]}$ is the subgroup of G of matrices with coefficients powers that are multiples of k .
- Decompose $d = p_1^{a_1} \dots p_r^{a_r}$, and let $\alpha_i = p_i^{a_i}, \beta_i = \frac{d}{\alpha_i}$.

Lemma

The $\overline{G}^{[\beta_i]}$ are p_i -Sylow of \overline{G} , they commute two by two and their product is \overline{G} .

($H, K < G$ commute means $HK = KH$, i.e. $\forall h, k, \exists h', k', hk = k'h'$.)

Reconstructing

This provides an alternative version of the Matched Product (Bachiller '18, Catino-Colazzo-Stefanelli '20) :

Suppose $(S, *_1)$, $(S, *_2)$ are cycle sets of size n and coprime classes d_1, d_2 . In $GL_n(\mathbb{C})$, consider $\overline{G} = \overline{G}_1 \overline{G}_2$.

Theorem (F.)

If $(S, *_1)$ and $(S, *_2)$ satisfy a "mixed" cycle set condition:

$$\forall s, t, u \in S, (s *_1 t) *_2 (s *_1 u) = (t *_2 s) *_1 (t *_2 u)$$

Then \overline{G} induces a cycle set structure on S of class (dividing) $d = d_1 d_2$.

We can restrict to cycle sets of class a prime-power to classify all cycle sets!

Indecomposability

Definition

$(S, *)$ is said to be decomposable if there exists a partition $S = X \sqcup Y$ such that $(X, *|_X), (Y, *|_Y)$ are cycle sets. Otherwise $(S, *)$ is called indecomposable.

- Up to a change of enumeration, S is decomposable iff the generators are diagonal by same blocks.

Proposition


If S is indecomposable and $d = p^k$, then $n = p^l$.

- We can "restrict" to cycle sets of size and class powers of the same prime.


Example: $(S, *)$ with $n = 16$ and $d = 30$ splits as cycle sets of class 2, 3 and 5. Those with class 3 and 5 must be decomposable.

Thank you for your attention!

Some histograms



Some histograms



An example

Let $S = \{s_1, \dots, s_6\}$ with $\psi(s_i) = (12 \dots 6) = \sigma$. Then:

- $s_i * s_j = s_{\sigma(i)} \Rightarrow T = \sigma$
 - $\psi(s_{i_1} \dots s_{i_k}) = \sigma^k \Rightarrow d = 6$
 - $\mathcal{G} = \langle \sigma \rangle \simeq \mathbb{Z}/6\mathbb{Z}$
-
- $G^{[3]}$ is generated by the $s_i^{[3]} = D_i^3 P_{\sigma^3}$, where $\sigma^3 = (14)(25)(36)$.
 - $S^{[3]}$ is of class 2
 - $\overline{G} = \overline{G}^{[2]} \overline{G}^{[3]}$, $s_i = (s_i^{[2]})^2 \cdot s_{\sigma^4(i)}^{[3]}$

Reconstructing

Denote Σ_n^d the group of monomial matrices with coefficient powers of ζ_d .
Define $\iota_d^{dk} : \Sigma_n^d \hookrightarrow \Sigma_n^{dk}$ sending ζ_d to ζ_{dk}^k .

- Let $(S, *_1)$ and $(S, *_2)$ be two cycle set structure on S .
- Suppose their classes d_1 and d_2 are coprime.
- Denote $\overline{G}_i < \Sigma_n^{d_i}$ their germs.
- Let $d = d_1 d_2$ and $\overline{G} = \iota_{d_1}^d(\overline{G}_1) \iota_{d_2}^d(\overline{G}_2)$.
- Bézout $\Rightarrow \exists u, v \in \mathbb{N}, d_2 u + d_1 v = 1[d] \Rightarrow \forall s \in S, \exists g \in \overline{G}, D_g = D_s$.

Does \overline{G} induces a cycle set structure on S ?

- No in general. Yes if \overline{G}_1 and \overline{G}_2 commute in Σ_n^d !

Example

$S = \{s_1, \dots, s_6\}$ with $(S', *_1)$ et $(S'', *_2)$ given by:

$$\psi_1(\{s'_1, \dots, s'_6\}) = (14)(25)(36) \quad d_1 = 2$$

$$\psi_2(\{s''_1, s''_3, s''_5\}) = (135) \quad \psi_2(\{s''_2, s''_4, s''_6\}) = (246) \quad d_2 = 3$$

• $3u + 2v = 1[6] \Rightarrow u = 1, v = 2$

$$\bar{s}_i = \iota_2^6(\overline{s'_i}^{[u]}) \iota_3^6(\overline{s''_{\psi(s'_i)(i)}}^{[v]})$$

• $\iota_2^6(\overline{s'_1}^{[1]}) = \iota_2^6 \begin{pmatrix} 000 \zeta_2 00 \\ 000 0 10 \\ 000 0 01 \\ 100 0 00 \\ 010 0 00 \\ 001 0 00 \end{pmatrix} = \begin{pmatrix} 000 \zeta_6^3 00 \\ 000 0 10 \\ 000 0 01 \\ 100 0 00 \\ 010 0 00 \\ 001 0 00 \end{pmatrix}$

• $\iota_3^6(\overline{s''_4}^{[2]}) = \begin{pmatrix} 1 & 0 & 0000 \\ 0 & 0 & 0001 \\ 0 & 0 & 1000 \\ 0 & (\zeta_6^2)^2 & 0000 \\ 0 & 0 & 0010 \\ 0 & 0 & 0100 \end{pmatrix}$

$$\bar{s}_1 = \begin{pmatrix} 0 \zeta_6 0000 \\ 0 0 0010 \\ 0 0 0100 \\ 1 0 0000 \\ 0 0 0001 \\ 0 0 1000 \end{pmatrix} \Rightarrow \text{We take } \psi(s_1) = (125634)$$

• We find: $\psi(\{s_1, s_3, s_5\}) = (125634)$, $\psi(\{s_2, s_4, s_6\}) = (145236)$